

RESH START TRAINING: BRIEF INTERVENTIONS WITH SMOKERS SMOKING KILLS

The size of the problem:

- · 21% of British people smoke (ONS, 2007)
- Smoking is by far the biggest single preventable cause of death and disease in the UK. One in two long-term smokers will die prematurely as a result of smoking, half of these in middle age (ASH, 2008)
- An estimated 110,000 deaths in the UK can be attributed to smoking annually - this is about one fifth of all UK deaths (Allender, 2009)
- The NHS currently spends more than £5 billion each year treating smoking-related diseases (Allender, 2009)

BUT...

70% of smokers say they want to give up (ONS, 2007)

Local Facts

RESH START TRAINING: BRIEF INTERVENTIONS WITH SMOKERS

- Derby City's smoking prevalence is 25%
- Highest prevalence is in Sinfin = 40%
- In 2009 / 10 Fresh Start supported ** to quit smoking
- In 2009 / 10 4043 people set a quit date
- Of the 4043 people who set quit date 12% (476) of these came from BME communities
- Of the 476 who set quit date 49% quit smoking

SUPPORT – FRESH START

The local NHS stop smoking service for Derby:

- Weekly advice and support for up to 12 weeks
- Up to 12 weeks' NRT on a voucher scheme, or other medication on prescription
- Choice of group, drop-in, one-to-one or telephone support Associate Advisors offer services in the community (Pharmacies, GP Surgeries, etc) as part of the care they provide
- Success rates of ~60% at four weeks
- Ongoing (52 week) support from Specialist Service

FRESH START IN THE COMMUNITY

· Specialist areas :

Acute trust Pregnancy

- GPs / Pharmacy Mental Health BME communities
- Routine and Manual workforces Young people Training

BME Communities & Young People Support for BME smokers: Clinics

Use of interpreters

FRESH START TRAINING: BRIEF INTERVENTIONS WITH SMOKERS

- Work with EMAC
- Ramadam campaign
- Local pharmacy / GP delivery

Support for Young People

- Dedicated YP worker
- Trained school nurses
- Clinics
- Promotion

FRESH START TRAINING BRIEF INTERVENTIONS WITH SMOKERS

• Secondhand smoke or 'passive smoking' is other people's tobacco smoke in the atmosphere. It is a major cause of indoor air pollution.

 85% of SHS is invisible. It can linger for at least two hours, even with a window open

•42% of children live in a home where at least one person smokes

•17000 under five year olds are hospitalised every year due to the effects of SHS

•Children at greater risk as immune systems less developed & have smaller airways

 Increased risk of: cot death, ear infections, respiratory illness & asthma





